

Lagrangian approach to local symmetries and self-dual model in gauge invariant formulation

Yong-Wan Kim* and Klaus D. Rothe†

Institut für Theoretische Physik

Universität Heidelberg

Philosophenweg 16, D-69120 Heidelberg

Abstract

Taking the Stückelberg Lagrangian associated with the abelian self-dual model of P. K. Townsend et al as a starting point, we embed this mixed first- and second-class system into a pure first-class system by following systematically the generalized Hamiltonian approach of Batalin, Fradkin and Tyutin. The resulting Lagrangian possesses an extended gauge invariance and provides a non-trivial example for a general Lagrangian approach to unravelling the full set of local symmetries of a Lagrangian.

PACS: 11.10.Ef, 11.10.Kk, 11.15.-q

Keywords: Hamiltonian embedding; Second-class; Self-dual;

Local symmetries; Lagrangian approach;

*e-mail: ywkim@physics.sogang.ac.kr, Department of Physics and Basic Science Research Institute, Sogang University, C.P.O.Box 1142, Seoul 100-611, Korea

†e-mail: k.rothe@thphys.uni-heidelberg.de

1 Introduction

The Hamiltonian approach to the quantization of constrained systems possessing first-class constraints has the drawback of not necessarily leading to a manifestly Lorentz covariant partition function. This problem is avoided in the Lagrangian field-antifield approach [1], which is based on an analysis of the local symmetries of a Lagrangian. The full set of such symmetries may not be evident from the outset, if the Lagrangian is of a more complicated nature. Their systematic and exhaustive determination thus constitute an integral part of the field-antifield quantization program.

In this paper, we reconsider the abelian version of the self-dual (SD) model of Townsend et al [2]. This model corresponds to a purely second-class system in the terminology of Dirac [3]. Following the generalized Hamiltonian approach of Batalin, Fradkin and Tyutin (BFT) [4], Kim et al [5] have turned all of the second-class constraints into first-class ones[‡]. The resulting Lagrangian was found to be given as the sum of a Stückelberg type Lagrangian [7] and a “Wess-Zumino” Lagrangian lacking manifest Lorentz, and even rotational invariance.

On the other hand, the non-abelian SD-model has been shown to be equivalently described by a Stückelberg Lagrangian involving mixed first- and second-class constraints [8]. Taking the abelian restriction of this Lagrangian as a starting point we obtain, following the systematic procedure of BFT, an equivalent Lagrangian involving only first-class constraints, exhibiting manifest rotational invariance. The equivalence with the formulation of [5] is established after suitable integration over one of the auxiliary fields in the extended phase space. This is done in Sections 2 and 3.

In Section 4, we then take the resulting first-class Lagrangian of Section 3 and systematically unravel all of its local symmetries by following a method

[‡]For a discussion following the Batalin-Fradkin approach see [6]

recently discussed in the literature [9]. Although these symmetries could of course be derived from the knowledge of the first-class constraints of the BFT Hamiltonian construction, it turns out that the model provides an interesting application of the Lagrangian approach of ref. [9], going well beyond the examples discussed in that reference.

We conclude in Section 5 with a summary.

2 First Class Formulation of Self-Dual Model

Our starting point is the Lagrangian of the abelian self-dual model of Townsend et al. [2],

$$\mathcal{L}_{SD} = \frac{1}{2}m^2 B^\mu B_\mu - \frac{m}{2}\epsilon_{\mu\nu\rho} B^\mu \partial^\nu B^\rho. \quad (2.1)$$

This Lagrangian describes a purely second-class system. As was shown in ref. [8], following the procedure of BFT for Hamiltonian embedding, this Lagrangian turns into a Lagrangian of the Stückelberg type, if one pair of second-class constraints is turned into first-class constraints. The result of this is a Lagrangian in which the fields B^μ have simply been gauged,

$$\tilde{\mathcal{L}} = \frac{1}{2}m^2(B_\mu + \partial_\mu\theta)(B^\mu + \partial^\mu\theta) - \frac{m}{2}\epsilon_{\mu\nu\rho} B^\mu \partial^\nu B^\rho. \quad (2.2)$$

The Lagrangian $\tilde{\mathcal{L}}$ describes a mixed system of first- and second-class constraints. We shall take it as a starting point for applying the BFT method [4] to also convert the remaining second-class constraints to first-class ones.

From (2.2) we obtain for the primary constraints,

$$\begin{aligned} T_0 &:= \pi_0 = 0, \\ T_i &:= \pi_i + \frac{1}{2m}\epsilon_{ik}B^k = 0. \end{aligned}$$

The canonical Hamiltonian associated with (2.2) is given by

$$H_c = \int d^2y \left\{ \frac{1}{2}\pi_\theta^2 + \frac{1}{2}B_i^2 - B^i \partial_i \theta + \frac{1}{2}(\partial_i \theta)^2 + B^0 (-\partial^i T_i + \frac{1}{m}\epsilon_{ij}\partial^i B^j - \pi_\theta) \right\}, \quad (2.3)$$

where

$$\pi_\theta = \partial^0 \theta + B^0. \quad (2.4)$$

The Dirac algorithm leads to a secondary constraint

$$T_3 := \partial^i T_i - \frac{1}{m} \epsilon_{ij} \partial^i B^j + \pi_\theta = 0. \quad (2.5)$$

The constraints T_0 and T_3 are seen to be first-class

$$\begin{aligned} \{T_0(x), T_\alpha(y)\} &= 0, \\ \{T_3(x), T_\alpha(y)\} &= 0, \quad \alpha = 0, 1, 2, 3, \end{aligned} \quad (2.6)$$

whereas the constraints $T_i = 0$ are found to be second class.

$$\{T_i(x), T_j(y)\} = \frac{1}{m} \epsilon_{ij} \delta^2(x - y) \equiv \Delta_{ij}(x, y). \quad (2.7)$$

The conversion of the second-class constraints T_i to first-class ones, Ω_i , follows the standard BFT procedure. For the constraints T_i one introduces “gauge-degrees of freedom” ϕ^i , $i = 1, 2$, together with the symplectic structure

$$\{\phi^i(x), \phi^j(y)\} = \omega^{ij}(x, y) \quad (2.8)$$

with ω^{ij} an arbitrary antisymmetric “matrix”, which we choose as

$$\omega^{ij}(x, y) = -\epsilon^{ij} \delta^2(x - y). \quad (2.9)$$

For the first-class constraints $\Omega_i = 0$, we make the ansatz

$$\Omega_i = \sum_{n=0}^{\infty} \Omega_i^{(n)}, \quad \Omega_i^{(0)} = T_i, \quad (2.10)$$

where $\Omega_i^{(n)}$ is a polynomial of degree n in the new fields ϕ^j . With the ansatz,

$$\Omega_i^{(1)}(x) = \int d^2 y X_{ij}(x, y) \phi^j(y), \quad (2.11)$$

the requirement of strong involution,

$$\{\Omega_i(x), \Omega_j(y)\} = 0, \quad (2.12)$$

then leads to recursive relations, plus the requirement

$$\Delta_{ij}(x, y) + \int d^2x' d^2y' X_{ik}(x, x') \omega^{kl}(x', y') X_{jl}(y', y) = 0. \quad (2.13)$$

This leads to the solution

$$X_{ij}(x, y) = \frac{1}{\sqrt{m}} \delta_{ij} \delta^2(x - y) \quad (2.14)$$

with the inverse

$$X^{ij}(x, y) = \sqrt{m} \delta_{ij} \delta^2(x - y), \quad (2.15)$$

as well as to the constraints $\Omega_\alpha = 0$, with

$$\begin{aligned} \Omega_0 &= \pi_0, \\ \Omega_i &= \pi_i + \frac{1}{2m} \epsilon_{ij} B^j + \frac{1}{\sqrt{m}} \phi^i, \\ \Omega_3 &= \partial^i \Omega_i^{(0)} - \frac{1}{m} \epsilon_{ij} \partial^i B^j + \pi_\theta. \end{aligned} \quad (2.16)$$

For the first-class Hamiltonian we similarly make the ansatz

$$\tilde{H} = H_c + \sum_{n=1}^{\infty} H^{(n)}. \quad (2.17)$$

The requirement that $\tilde{H}$ be in strong involution with the first-class constraints then leads to the recursive relations

$$H^{(n)} = -\frac{1}{n} \int d^2x d^2y d^2z \phi^i(x) \omega_{ij}(x, y) X^{jk}(y, z) G_k^{(n-1)}(z), \quad (n \geq 1), \quad (2.18)$$

where the generating functionals $G_k^{(n)}$ are given by

$$\begin{aligned} G_i^{(0)} &= \{\Omega_i^{(0)}, H_c\}_O, \\ G_i^{(n)} &= \{\Omega_i^{(0)}, H^{(n)}\}_O + \{\Omega_i^{(1)}, H^{(n-1)}\}_O, \end{aligned} \quad (2.19)$$

where the subscript O means that Poisson brackets are to be calculated with respect to the original variables. We find recursively

$$\begin{aligned} G_i^{(0)} &= B_i + \partial_i \theta, \quad H^{(1)} = \sqrt{m} \int d^2x \phi^i \epsilon_{ij} (B^j + \partial^j \theta), \\ G_i^{(1)} &= \sqrt{m} \epsilon_{ij} \phi^j, \quad H^{(2)} = \frac{m}{2} \int d^2x \phi^i \phi^i, \\ G_i^{(n)} &= 0, \quad H^{(n+1)} = 0, \quad n \geq 2. \end{aligned} \quad (2.20)$$

One readily checks that the final Hamiltonian

$$\tilde{H} = H_c + H_1 + H_2 \quad (2.21)$$

is indeed strongly involutive with respect to the constraints.

3 The Partition Function

The partition function corresponding to the Hamiltonian (2.21) reads

$$Z = \int D\pi_\mu DB^\mu \int D\pi_\theta D\theta \int D\phi^1 D\phi^2 \prod_\alpha \delta[\Omega_\alpha] \prod_\beta \delta[\Gamma_\beta] \det\{\Omega_\alpha, \Gamma_\beta\} e^{iS'}, \quad (3.1)$$

where

$$S' = \int d^3x \left\{ \pi_\mu \dot{B}^\mu + \pi_\theta \dot{\theta} + \frac{1}{2} \phi^i \epsilon_{ij} \dot{\phi}^j - \mathcal{H}_c - \sqrt{m} \phi^i \epsilon_{ij} (B^j + \partial^j \theta) - \frac{m}{2} \phi^i \phi^i \right\}, \quad (3.2)$$

and Γ_β are gauge-fixing conditions. Noting that

$$\prod_i \delta[\Omega_i] \delta[\Omega_3] = \prod_i \delta[\Omega_i] \delta\left[-\frac{1}{m} \epsilon_{jk} \partial^j B^k + \pi_\rho - \frac{1}{\sqrt{m}} \partial^j \phi^k\right], \quad (3.3)$$

and writing the second δ -function as a Fourier integral with Fourier variable ξ , we obtain, after performing the π_μ integration and making the shift of variable $B^0 + \xi \rightarrow B^0$,

$$Z = \int D\xi \int DB^\mu \int D\pi_\theta D\theta \int \prod_{i=1}^2 D\phi^i \prod_\beta \delta[\Gamma_\beta] \det\{\Omega_\alpha, \Gamma_\beta\} e^{iS''}, \quad (3.4)$$

where

$$S'' = S + S_{WZ} \quad (3.5)$$

with S the action corresponding to the Lagrangian (2.2), and

$$S_{WZ} = \int d^2x \left\{ -\frac{1}{\sqrt{m}} \phi^i B^{0i} - \sqrt{m} \phi^i \epsilon_{ij} (B^j + \partial^j \theta) + \frac{1}{2} \phi^i \epsilon_{ij} \partial_0 \phi^j - \frac{m}{2} \phi^i \phi^i \right\}. \quad (3.6)$$

Unlike the action (51) of ref. [5], the action (3.5) is manifestly rotationally invariant. Nevertheless, the corresponding partition functions are equivalent.

Indeed, integration over ϕ^1 in (3.4) turns Z into the partition function of ref. [5].

To conclude this section, let us show that unlike in the case of the configuration space partition function of ref. [5], there exists a “canonical” gauge in which we recover the partition function of the Stückelberg formulation.

In the Hamiltonian formulation, the first-class constraints (2.16) are generators of local symmetry transformations. Thus, defining

$$G = \sum_{\alpha} \int d^2x \epsilon^{\alpha}(x) \Omega_{\alpha}(x), \quad (3.7)$$

we have ($\delta A = \{A, g\}$)

$$\begin{aligned} \delta B^0 &= \epsilon^0, \\ \delta B^i &= \epsilon^i + \partial^i \epsilon^3, \\ \delta \theta &= \epsilon^3, \\ \delta \phi^i &= -\frac{1}{\sqrt{m}} \epsilon_{ij} \epsilon^j. \end{aligned} \quad (3.8)$$

One easily checks that the transformation (3.8) leaves the action (3.5) invariant, provided that we choose $\epsilon^0 = \partial_0 \epsilon^3$ [§]. Consider in particular the transformation with $\epsilon^0 = \epsilon^3 = 0$, and $\epsilon^j = \sqrt{m} \epsilon_{jk} \phi^k$. Under this transformation $\phi^i \rightarrow 0$, $i = 1, 2$; the gauge $\phi^1 = \phi^2 = 0$ is thus accessible, and the partition function (3.4) reduces in this gauge to that of the Stückelberg formulation (2.2).

4 Lagrangian Approach to Local Symmetries

In the field-antifield formalism [1] the establishment of the full, irreducible set of local symmetries of a Lagrangian plays a fundamental role. In this formalism these symmetries must be identified without the use of Dirac’s

[§]As is well known, symmetry transformations of the Lagrangian generally imply restrictions on the symmetry transformations of the total Hamiltonian.

Hamiltonian construction of the corresponding generators. In this section we illustrate a recently proposed [9] Lagrangian approach in terms of the action (3.5) in the extended configuration space. Our notation deviates somewhat from that of ref. [9].

The equations of motion following from (3.5) are of the form

$$L_{ix} = \int d^3y (W_{ix,jy} \ddot{\varphi}_{jy} + \alpha_{ix}) = 0, \quad i = 1, 2, \dots, 6, \quad (4.1)$$

where

$$(\varphi_{ix})^T = (B^0(x), B^1(x), B^2(x), \phi^1(x), \phi^2(x), \theta(x)), \quad (4.2)$$

$W_{ix,jy}$ is the Hessian

$$\begin{aligned} W_{ix,jy} &= \frac{\delta^2 L}{\delta \dot{\varphi}_{ix} \delta \dot{\varphi}_{jy}} = \delta_{i6} \delta_{j6} \delta^3(x - y) \\ &\equiv \tilde{W}_{ij} \delta^3(x - y), \end{aligned} \quad (4.3)$$

and

$$(\alpha_{ix})^T = (\alpha_{B^0}(x), \alpha_{B^1}(x), \alpha_{B^2}(x), \alpha_{\phi^1}(x), \alpha_{\phi^2}(x), \alpha_{\theta}(x)) \quad (4.4)$$

with

$$\begin{aligned} \alpha_{B^0} &= \frac{1}{m} \epsilon_{ij} \partial^i B^j - (B^0 + \partial^0 \theta) + \frac{1}{\sqrt{m}} \partial^i \phi^i, \\ \alpha_{B^i} &= -\frac{1}{m} \epsilon_{ij} (\partial^0 B^j - \partial^j B^0) + (B^i + \partial^i \theta) - \frac{1}{\sqrt{m}} \partial^0 \phi^i - \sqrt{m} \epsilon_{ij} \phi^j, \\ \alpha_{\phi^i} &= -\epsilon_{ij} \partial_0 \phi^j + \frac{1}{\sqrt{m}} (\partial^0 B^i - \partial^i B^0) + \sqrt{m} \epsilon_{ij} (B^j + \partial^j \theta) + m \phi^i, \\ \alpha_{\theta} &= \partial^0 B^0 + \partial_i (B^i + \partial^i \theta) + \sqrt{m} \epsilon_{ij} \partial^i \phi^j. \end{aligned} \quad (4.5)$$

The Hessian matrix $\tilde{W}_{ij}$ (4.3) is of rank one. Hence there exist five “zeroth generation” null eigenvectors (λ_{ix}) satisfying

$$\int d^3x \lambda_{ix} W_{ix,jy} = 0. \quad (4.6)$$

We choose them to have components

$$\lambda_{ix}(a, z) = \delta^3(x - z) \delta_{ia}, \quad a = 1, \dots, 5, \quad (4.7)$$

where (a, z) label the eigenvectors and (i, x) label the components.

Correspondingly we have the “zeroth generation” constraints

$$\Omega_a(z) = \int d^3x \lambda_{ix}(a, z) L_{ix} = \alpha_a(z) = 0, \quad a = 1, \dots, 5. \quad (4.8)$$

From here on we shall drop continuum labels, which are now implicitly contained in the associated discrete labels. From (4.4) we see that only three of these represent non-trivial constraints, since we have identically

$$\alpha_{B^i} + \frac{1}{\sqrt{m}} \epsilon_{ij} \alpha_{\phi^j} \equiv 0. \quad (4.9)$$

Introducing the null eigenvectors

$$\begin{aligned} \hat{\lambda}(1) &= (0, 1, 0, 0, \frac{1}{\sqrt{m}}, 0), \\ \hat{\lambda}(2) &= (0, 0, 1, \frac{-1}{\sqrt{m}}, 0, 0), \end{aligned} \quad (4.10)$$

we can write the identities in the form

$$\hat{\Omega}_{\hat{a}} := \hat{\lambda}_i(\hat{a}) L_i \equiv 0, \quad \hat{a} = 1, 2. \quad (4.11)$$

For the remaining, non-trivial constraints we choose

$$\bar{\Omega}_{\bar{a}} := \bar{\lambda}_i(\bar{a}) L_i = 0, \quad \bar{a} = 1, 2, 3 \quad (4.12)$$

with

$$\bar{\lambda}_i(\bar{a}) = \lambda_i(a). \quad (4.13)$$

This also establishes our notation: quantities related to trivial (non-trivial) constraints are denoted by a “hat” (“bar”).

We now require the (non-trivial) constraints (4.12) to be independent of time. We thus need to add to the equations of motion (4.1) the equations $\dot{\bar{\Omega}}_{\bar{a}} = 0$, $\bar{a} = 1, 2, 3$. The resulting set of 9 equations may be summarized in the form of the set of “first generation” equations $L_{i_1}^{(1)} = 0$, $i_1 = 1, \dots, 9$, with

$$L_{i_1}^{(1)} = \begin{cases} L_i, & i = 1, \dots, 6, \\ \frac{d}{dt}(\bar{\lambda}_i(\bar{a}) L_i), & i = 6 + \bar{a}, \quad \bar{a} = 1, 2, 3. \end{cases} \quad (4.14)$$

$L_{i_1}^{(1)}$ can be written in the general form

$$L_{i_1}^{(1)} := W_{i_1 j}^{(1)} \ddot{\phi}_j + \alpha_{i_1}^{(1)} = 0, \quad (4.15)$$

where

$$(W_{i_1 j}^{(1)}) = \left(\begin{array}{c} (\tilde{W}_{ij}) \\ \hline \begin{array}{cccccc} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & \frac{1}{m} & \frac{1}{\sqrt{m}} & 0 & 0 \\ 0 & -\frac{1}{m} & 0 & 0 & \frac{1}{\sqrt{m}} & 0 \end{array} \end{array} \right) \delta^2(x-y), \quad (4.16)$$

and

$$(\alpha_{i_1}^{(1)}) = \left(\begin{array}{c} (\alpha_i) \\ \hline \begin{array}{c} \alpha_7^{(1)} \\ \alpha_8^{(1)} \\ \alpha_9^{(1)} \end{array} \end{array} \right) \quad (4.17)$$

with

$$\begin{aligned} \alpha_7^{(1)} &= -\frac{1}{m} \partial^1 \dot{B}^2 + \frac{1}{m} \partial^2 \dot{B}^1 + \dot{B}^0 - \frac{1}{\sqrt{m}} \partial^i \dot{\phi}^i, \\ \alpha_8^{(1)} &= -\frac{1}{m} \partial^2 \dot{B}^0 - (\dot{B}^1 + \partial^1 \dot{\theta}) + \sqrt{m} \dot{\phi}^2, \\ \alpha_9^{(1)} &= \frac{1}{m} \partial^1 \dot{B}^0 - (\dot{B}^2 + \partial^2 \dot{\theta}) - \sqrt{m} \dot{\phi}^1, \end{aligned} \quad (4.18)$$

respectively.

We now repeat the previous analysis taking eqns. (4.15) as a starting point, and looking for solutions of

$$\lambda_{i_1}^{(1)}(a_1)W_{i_1 j}^{(1)} = 0. \quad (4.19)$$

The null eigenvectors of $W^{(1)}$ are of the form $(\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6, -\lambda_4, 0, 0)$. In addition to $(\vec{\lambda}(\bar{a}), 0, 0, 0)$, $\bar{a} = 1, 2, 3$ and $(\vec{\lambda}(\hat{a}), 0, 0, 0)$, $\hat{a} = 1, 2$, we have one further null eigenvector, which we choose as ¶

$$\lambda^{(1)}(1) = (0, 0, 0, 0, 0, 1, -1, 0, 0). \quad (4.20)$$

Associated with this eigenvector we have a new constraint

$$\Omega_1^{(1)} := \lambda_{i_1}^{(1)} \alpha_{i_1}^{(1)} = 0. \quad (4.21)$$

Explicitly

$$\begin{aligned} \Omega_1^{(1)} &= \alpha_6^{(1)} - \alpha_7^{(1)} \\ &= \partial_i(B^i + \partial^i \theta) + \sqrt{m} \epsilon_{ij} \partial^i \phi^j + \frac{1}{m} \epsilon_{ij} \partial^i \dot{B}_j + \frac{1}{\sqrt{m}} \partial^i \dot{\phi}^i. \end{aligned} \quad (4.22)$$

Since this represents a non-trivial constraint, we write $\Omega_1^{(1)} = \bar{\Omega}_1^{(1)}$, $\lambda^{(1)}(1) = \bar{\lambda}^{(1)}(\bar{1})$. Correspondingly (4.21) may be written in the form

$$\bar{\Omega}_{\bar{a}_1}^{(1)} = \bar{\lambda}_{i_1}^{(1)}(\bar{a}_1) L_{i_1}^{(1)} = 0, \quad \bar{a}_1 = 1. \quad (4.23)$$

There exist no new identities at this level.

We repeat the procedure by requiring $\dot{\bar{\Omega}}_{\bar{a}_1}^{(1)} = 0$, i.e.

$$\frac{1}{\sqrt{m}} \partial^i \ddot{\phi}^i + \frac{1}{m} \epsilon_{ij} \partial^i \ddot{B}^j + \partial_i(\dot{B}^i + \partial^i \dot{\theta}) + \sqrt{m} \epsilon_{ij} \partial^i \dot{\phi}^j = 0. \quad (4.24)$$

Adjoining this equation to $L_{i_1}^{(1)} = 0$, we have $L_{i_2}^{(2)} = 0$, $i_2 = 1, \dots, 10$, with

$$L_{i_2}^{(2)} = \begin{cases} L_i, & i = 1, \dots, 6, \\ \frac{d}{dt}(\bar{\lambda}_i(\bar{a}) L_i), & i_2 = 6 + \bar{a}, \bar{a} = 1, 2, 3, \\ \frac{d}{dt}(\bar{\lambda}_{i_1}^{(1)}(\bar{a}_1) L_{i_1}^{(1)}), & i_2 = 9 + \bar{a}_1, \bar{a}_1 = 1. \end{cases} \quad (4.25)$$

¶Each component is understood to be multiplied by $\delta^2(x - z)$.

The resulting complete set of equations is now of the form

$$L_{i_2}^{(2)} := W_{i_2 j}^{(2)} \ddot{\varphi}_j + \alpha_{i_2}^{(2)} = 0, \quad i_2 = 1, \dots, 10, \quad (4.26)$$

where

$$\begin{aligned} W_{i_2 j}^{(2)} &= W_{i_2 j}, \quad i_2 = 1, \dots, 6, \\ W_{i_2 j}^{(2)} &= W_{i_2 j}^{(1)}, \quad i_2 = 7, 8, 9, \\ W_{10, j}^{(2)} &= (0, -\frac{\partial^2}{m}, \frac{\partial^1}{m}, \frac{\partial^1}{\sqrt{m}}, \frac{\partial^2}{\sqrt{m}}, 0) \delta^2(x - y), \end{aligned} \quad (4.27)$$

and

$$\begin{aligned} \alpha_{i_2}^{(2)} &= \alpha_{i_2}, \quad i_2 = 1, \dots, 6, \\ \alpha_{i_2}^{(2)} &= \alpha_{i_2}^{(1)}, \quad i_2 = 7, 8, 9, \\ \alpha_{10}^{(2)} &= \partial_i (\dot{B}^i + \partial^i \dot{\theta}) + \sqrt{m} \epsilon_{ij} \partial^i \dot{\phi}^j. \end{aligned} \quad (4.28)$$

In addition to the previous null eigenvectors (enhanced by a suitable number of zeroes in the last components), we thus have the new null eigenvector

$$\lambda^{(2)}(1) = (0, 0, 0, 0, 0, 0, 0, \partial_x^1, \partial_x^2, -1) \delta^2(x - z). \quad (4.29)$$

The associated constraint is found to vanish “identically”:

$$\lambda_{i_2}^{(2)}(1) \alpha_{i_2}^{(2)} = \partial^1 \alpha_8^{(2)} + \partial^2 \alpha_9^{(2)} - \alpha_{10}^{(2)} = 0. \quad (4.30)$$

Hence we denote $\lambda^{(2)}(1)$ by $\hat{\lambda}^{(2)}(\hat{1})$ and write (4.30) in the form

$$\hat{\Omega}_{\hat{a}_2}^{(2)} := \hat{\lambda}_{i_2}^{(2)}(\hat{a}_2) L_{i_2}^{(2)} \equiv 0, \quad \hat{a}_2 = 1. \quad (4.31)$$

The algorithm ends at this point.

The local symmetries of the action (3.5) are encoded in the identities (4.11) and (4.31). Recalling (4.14) and (4.25) we see that these identities can be rewritten as follows:

$$\begin{aligned} \hat{\Omega}_1 &= L_2 + \frac{1}{\sqrt{m}} L_5 \equiv 0, \\ \hat{\Omega}_2 &= L_3 - \frac{1}{\sqrt{m}} L_4 \equiv 0, \\ \hat{\Omega}_1^{(2)} &= \frac{d}{dt} (\partial^1 L_2 + \partial^2 L_3 + L_6) + \frac{d^2}{dt^2} L_1 \equiv 0. \end{aligned} \quad (4.32)$$

This result is a special case of a general theorem stating [9] that the identities

$\hat{\Omega}_{\hat{a}_l}^{(l)} \equiv 0$ ($\hat{\Omega}_{\hat{a}_0}^{(0)} \equiv \hat{\Omega}_{\hat{a}}$) can always be written in the form

$$\hat{\Omega}_{\hat{a}_l}^{(l)} = \sum_{s=0}^l \frac{d^s}{dt^s} \left(\Phi_{si}^{(l)}(\hat{a}_l) L_i \right). \quad (4.33)$$

For the case in question

$$\begin{aligned} \Phi_{0i}^{(0)}(1) &= \delta_{i2} + \frac{1}{\sqrt{m}} \delta_{i5}, \\ \Phi_{0i}^{(0)}(2) &= \delta_{i3} - \frac{1}{\sqrt{m}} \delta_{i4}, \\ \Phi_{1i}^{(1)}(1) &= \delta_{i2} \partial^1 + \delta_{i3} \partial^2 + \delta_{i6}, \\ \Phi_{0i}^{(2)}(1) &= 0 \\ \Phi_{2i}^{(2)}(1) &= \delta_{i1}. \end{aligned} \quad (4.34)$$

It again follows from general considerations [9] that the action (3.5) is invariant under the transformation

$$\delta \varphi_i = \sum_{l=0}^l \sum_{s=0}^l \left(\sum_{\hat{a}_l} (-1)^s \Phi_{si}^{(l)}(\hat{a}_l) \frac{d^s}{dt^s} \epsilon_{\hat{a}_l}^{(l)} \right). \quad (4.35)$$

For the case in question this corresponds to the transformations

$$\begin{aligned} \delta B^0 &= \ddot{\epsilon}_1^{(2)}, \\ \delta B^1 &= \epsilon_1^{(0)} - \partial^1 \dot{\epsilon}_1^{(1)}, \\ \delta B^2 &= \epsilon_2^{(0)} - \partial^2 \dot{\epsilon}_1^{(1)}, \\ \delta \phi^1 &= -\frac{1}{\sqrt{m}} \epsilon_2^{(0)}, \\ \delta \phi^2 &= \frac{1}{\sqrt{m}} \epsilon_1^{(0)}, \\ \delta \theta &= -\dot{\epsilon}_1^{(1)}. \end{aligned} \quad (4.36)$$

Comparison with (3.8) with $\epsilon^0 = \ddot{\epsilon}_1^{(2)}$, $\epsilon^3 = -\dot{\epsilon}_1^{(1)}$, and $\epsilon^i = \epsilon_i^{(0)}$ shows that we have recovered the local symmetry transformations.

5 Conclusion

In this paper we reconsidered the abelian self-dual model of Townsend et al [2], by taking as a starting point its description as a mixed first- and second-class system in terms of a Stückelberg type Lagrangian [7]. The equivalence of this description with the original, purely second-class system was established in [8] for the non-abelian case, and hence in particular for the abelian case. We turned this Stückelberg Lagrangian into a purely first-class system, following the method of Batalin, Fradkin and Tyutin [4]. We then showed that the Lagrangian thus obtained was equivalent to the Lagrangian eq. (51) of ref. [5]. This established that despite the lack of manifest rotational invariance of that Lagrangian, it defined a theory consistent with rotational invariance.

The Lagrangian we obtained in this paper exhibited a larger local symmetry than the original Stückelberg Lagrangian, reflecting the existence of twice as many first-class constraints, that is, generators of gauge transformations. We then showed, following the method of ref. [9], how these symmetries could be systematically derived on a purely Lagrangian level, without resorting to a Hamiltonian formulation. The Lagrangian approach employed in this derivation should be of much interest in the context of the field-antifield formalism. As it turned out, our formulation of the self-dual model as a purely first-class system provided a non-trivial example for such a systematic construction.

Acknowledgement

One of the authors (Y. W. Kim) would like to thank the Institut für Theoretische Physik for the kind hospitality, and the Korea Research Foundation for (1996) overseas fellowship which made this collaboration possible.

References

- [1] J. Gomis, J. Paris and S. Samuel, Phys. Rep. **259** (1995) 1.
- [2] P. K. Townsend, K. Pilch and P. van Nieuwenhuizen, Phys. Lett. **136B** (1984) 38; **137B** (1984) 443; A. Karlhede, U. Lindström, M. Roček and P. van Nieuwenhuizen, Phys. Lett. **186B** (1987) 96.
- [3] P. A. M. Dirac, Can. J. Math. **2** (1950) 129: “Lectures on Quantum Mechanics” (Yeshiva University, New York, 1964).
- [4] I. A. Batalin and E. S. Fradkin, Phys. Lett. **B180** (1986) 157; Nucl. Phys. **B279** (1987) 514; I. A. Batalin and I. V. Tyutin, Int. J. Mod. Phys. **A6** (1991) 3255.
- [5] Y. W. Kim, Y. J. Park, K. Y. Kim and Y. Kim, Phys. Rev. **D51** (1995) 2943.
- [6] R. Banerjee, H. J. Rothe and K. D. Rothe, Nucl. Phys. **B426** (1994) 129; R. Banerjee and H. J. Rothe, Nucl. Phys. **B447** (1995) 183; R. Banerjee, H. J. Rothe and K. D. Rothe, Phys. Rev. **D52** (1995) 3750; *ibid.*, Phys. Rev. **D55** (1997) 1.
- [7] T. Kunimasa and T. Goto, Prog. Theor. Phys. **37** (1967) 452.
- [8] Y.-W. Kim and K. D. Rothe, Nucl. Phys. B (in press).
- [9] A. Shirzad, “Gauge Symmetry in Lagrangian formulation and Schwinger Model”, IPM-97-205; For a general discussion, see also, E. C. G. Sudarshan and N. Mukunda, “Classical Dynamics: A Modern Perspective”, (John Wiley & Sons, 1974).